

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : <b>C11D 3/48, 1/62, 1/645, 3/20, A01N 33/12</b>		<b>A1</b>	(11) International Publication Number: <b>WO 00/49127</b>
			(43) International Publication Date: 24 August 2000 (24.08.00)
(21) International Application Number: <b>PCT/GB00/00464</b>			(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
(22) International Filing Date: 14 February 2000 (14.02.00)			
(30) Priority Data: 9903478.7 17 February 1999 (17.02.99) GB			
(71) Applicant (for all designated States except US): RECKITT & COLMAN INC. [US/US]; 1655 Valley Road, Wayne, NJ 07474 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): SMIALOWICZ, Dennis, Thomas [US/US]; 40 John Dow Avenue, Waldwick, NJ 07463 (US). CHEUNG, Tak, Wai [GB/US]; 8 Jenna Drive, Bridgewater, NJ 08807 (US).			
(74) Agents: DICKSON, Elizabeth, Anne; Reckitt Benckiser plc, Group Patents Department, Dansom Lane, Hull HU8 7DS (GB) et al.			Published With international search report.
(54) Title: <b>HARD SURFACE CLEANING AND DISINFECTING COMPOSITIONS</b>			
(57) Abstract  Hard surface cleaning and disinfecting compositions comprise a synergistic combination of a quaternary ammonium compound and an alkoxylated quaternary ammonium compound. The hard surface cleaning and disinfecting compositions provide excellent cleaning and disinfection of hard surfaces.			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**HARD SURFACE CLEANING AND DISINFECTING COMPOSITIONS**

5

The present invention relates to hard surface cleaning and disinfecting compositions which comprise a synergistic combination of a quaternary ammonium compound and an alkoxylated quaternary ammonium compound.

10

The prior art has suggested many aqueous compositions which are directed to provide a cleaning or disinfecting benefit to such hard surfaces. These compositions predominantly are aqueous preparations which include one or more deterative surfactants, one or more organic solvents and in minor amounts, conventional additives included enhance the attractiveness of the product, typically fragrances and coloring agents. Certain of these also include one or more constituents which provide a primary disinfecting benefit to the aqueous preparations.

15

20

While these known-art compositions may provide advantages, there is a continuing need in the art for such hard surface treatment compositions which include reduced amounts of active constituents, and which minimize or eliminate the amounts of organic solvents which need be present in such compositions.

25

It is yet a further object of the invention to provide a readily pourable and readily pumpable cleaning composition which features the benefits described above.

30

It is a further object of the invention to provide a process for cleaning or sanitization of hard surfaces, which process comprises the step of: providing the composition as outlined above, and  
5 applying an effective amount to a hard surface requiring such treatment.

These and other objects of the invention shall be more apparent from a reading of the specification and of the claims attached.

10 According to a first aspect of the invention there is provided a hard surface cleaning and disinfecting composition which comprises (preferably, consisting essentially of) the following constituents:

- 15 (a) a germicidal constituent which provides a primary sanitizing benefit, preferably a quaternary ammonium compound;
- (b) alkoxyated quaternary ammonium compound;
- (c) optionally, one or more deterative surfactants particularly selected from carboxylate,  
20 nonionic, cationic and amphoteric surfactants;
- (d) optionally, one or more organic solvents;
- (e) a major proportion of water.

The compositions described above may include one  
25 or more further conventional optional constituents such as: pH buffering agents, perfumes, perfume carriers, colorants, hydrotropes, germicides, fungicides, anti-oxidants, anti-corrosion agents, and the like.

30 Preferred compositions according to the invention are largely aqueous, and are readily pourable and pumpable when packaged from a manually operable pump,

such as a 'trigger spray' dispenser. The preferred compositions of the invention feature good cleaning, disinfection of hard surfaces and little or not buildup of residue on treated hard surfaces.

5       According to a second aspect of the invention, there is provided a hard surface cleaning and disinfecting composition which comprises (preferably, consisting essentially of) the following constituents:

10       (a) a germicidal constituent which provides a primary sanitizing benefit, preferably a quaternary ammonium compound;

(b) alkoxyated quaternary ammonium compound;

15       (c) one or more deterative surfactants particularly selected from carboxylate, nonionic, cationic and amphoteric surfactants;

(e) a major proportion of water;

20       characterized in that the composition is essentially free of (d) one or more organic solvents, such as water soluble alcohols, ethers, and glycol ethers. These compositions may include one or more further conventional optional constituents such as: pH buffering agents, perfumes, perfume carriers, colorants, hydrotropes, germicides, fungicides, anti-oxidants, anti-corrosion agents, and the like.

25       According to a third aspect of the invention, there is provided a hard surface cleaning and disinfecting composition which comprises (preferably, consisting essentially of) the following constituents:

30       (a) a germicidal constituent which provides a primary sanitizing benefit, preferably a quaternary ammonium compound;

- (b) alkoxyated quaternary ammonium compound;
- (d) one or more organic solvents;
- (e) a major proportion of water.

characterized in that the composition is essentially  
5 free of (c) one or more deterative surfactants  
particularly selected from carboxylate, nonionic,  
cationic and amphoteric surfactants;

The compositions may include one or more further  
conventional optional constituents such as: pH  
10 buffering agents, perfumes, perfume carriers,  
colorants, hydrotropes, germicides, fungicides, anti-  
oxidants, anti-corrosion agents, and the like.

In accordance with a fourth aspect of the  
invention first aspect of the invention there is  
15 provided a hard surface cleaning and disinfecting  
composition which comprises (preferably, consisting  
essentially of) the following constituents:

- (a) a germicidal constituent which provides a  
primary sanitizing benefit, preferably a  
20 quaternary ammonium compound;
- (b) alkoxyated quaternary ammonium compound;
- (e) a major proportion of water;

wherein the compositions are essentially free of (c)  
deterative surfactants, particularly carboxylate,  
25 nonionic, cationic and amphoteric surfactants, as well  
as being essentially free of (d) organic solvents.


The compositions described above may include one  
or more further conventional optional constituents  
such as: pH buffering agents, perfumes, perfume  
30 carriers, colorants, hydrotropes, germicides,  
fungicides, anti-oxidants, anti-corrosion agents, and  
the like.

The compositions described above may include one or more further conventional optional constituents such as: pH buffering agents, perfumes, perfume carriers, colorants, hydrotropes, germicides, fungicides, anti-oxidants, anti-corrosion agents, and the like.

The inventive compositions necessarily include (a) at least one germicidal constituent which provides a primary sanitizing benefit to the compositions.

Particularly preferred for use as the (a) germicidal constituent is at least one cationic surfactant which is found to provide a broad antibacterial or sanitizing function. Any cationic surfactant which satisfies these requirements may be used and are considered to be within the scope of the present invention, and mixtures of two or more cationic surface active agents, viz., cationic surfactants may also be used. Cationic surfactants are well known, and useful cationic surfactants may be one or more of those described for example in *McCutcheon's Detergents and Emulsifiers*, North American Edition, 1982; *Kirk-Othmer, Encyclopedia of Chemical Technology*, 3rd Ed., Vol. 22, pp. 346-387, the contents of which are herein incorporated by reference.

Examples of preferred cationic surfactant compositions useful in the practice of the instant invention are those which provide a germicidal effect to the concentrate compositions, and especially preferred are quaternary ammonium compounds and salts thereof, which may be characterized by the general structural formula:



where at least one of  $R_1$ ,  $R_2$ ,  $R_3$  and  $R_4$  is a alkyl, aryl or alkylaryl substituent of from 6 to 26 carbon atoms, and the entire cation portion of the molecule

5 has a molecular weight of at least 165. The alkyl substituents may be long-chain alkyl, long-chain alkoxyaryl, long-chain alkylaryl, halogen-substituted long-chain alkylaryl, long-chain alkylphenoxyalkyl, arylalkyl, etc. The remaining substituents on the

10 nitrogen atoms other than the abovementioned alkyl substituents are hydrocarbons usually containing no more than 12 carbon atoms. The substituents  $R_1$ ,  $R_2$ ,  $R_3$  and  $R_4$  may be straight-chained or may be branched, but are preferably straight-chained, and may include one

15 or more amide, ether or ester linkages. The counterion X may be any salt-forming anion which permits water solubility of the quaternary ammonium complex.

Exemplary quaternary ammonium salts within the


20 above description include the alkyl ammonium halides such as cetyl trimethyl ammonium bromide, alkyl aryl ammonium halides such as octadecyl dimethyl benzyl ammonium bromide, N-alkyl pyridinium halides such as N-cetyl pyridinium bromide, and the like. Other

25 suitable types of quaternary ammonium salts include those in which the molecule contains either amide, ether or ester linkages such as octyl phenoxy ethoxy ethyl dimethyl benzyl ammonium chloride, N-(laurylcocoaminoformylmethyl)-pyridinium chloride, and



the like. Other very effective types of quaternary ammonium compounds which are useful as germicides include those in which the hydrophobic radical is characterized by a substituted aromatic nucleus as in  
 5 the case of lauryloxyphenyltrimethyl ammonium chloride, cetylaminophenyltrimethyl ammonium methosulfate, dodecylphenyltrimethyl ammonium methosulfate, dodecylbenzyltrimethyl ammonium chloride, chlorinated dodecylbenzyltrimethyl ammonium  
 10 chloride, and the like.

Preferred quaternary ammonium compounds which act as germicides and which are to be found useful in the practice of the present invention include those which have the structural formula:



15

wherein  $\text{R}_2$  and  $\text{R}_3$  are the same or different  $\text{C}_8$ - $\text{C}_{12}$ alkyl, or  $\text{R}_2$  is  $\text{C}_{12-16}$ alkyl,  $\text{C}_8$ - $\text{C}_{18}$ alkylethoxy,  $\text{C}_8$ - $\text{C}_{18}$ alkylphenoethoxy and  $\text{R}_3$  is benzyl, and X is a  
 20 halide, for example chloride, bromide or iodide, or is a methosulfate anion. The alkyl groups recited in  $\text{R}_2$  and  $\text{R}_3$  may be straight-chained or branched, but are preferably substantially linear.

Particularly useful quaternary germicides include  
 25 compositions which include a single quaternary compound, as well as mixtures of two or more different quaternary compounds. Particularly useful quaternary germicides include which are described as being a blend of alkyl dimethyl benzyl ammonium chlorides;


BARDAC® 205M, BARDAC® 2050, BARDAC® 2080, BARDAC® 2250, BTC® 812, BTC® 818 and BTC® 1010 which are described as being based on dialkyl(C<sub>8</sub>-C<sub>10</sub>)dimethyl ammonium chloride; BARDAC® 2250 and BARDAC® 2280 or  
5 BTC® 1010 which are described as being a composition which includes didecyl dimethyl ammonium chloride; BARDAC® LF and BARDAC® LF 80 which are described as being based on dioctyl dimethyl ammonium chloride; BARQUAT® MB-50, BARQUAT® MB-80, BARQUAT® MX-50,  
10 BARQUAT® MX-80, BARQUAT® OJ-50, BARQUAT® OJ-80, BARDAC® 208M, HYAMINE® 3500, HYAMINE® 3500-NF, BTC® 50, BTC® 824, BTC® 835, BTC® 885, BTC® 2565, BTC® 2658, BTC® 8248 or BTC® 8358 each described as being based on alkyl dimethyl benzyl ammonium chloride  
15 (benzalkonium chloride); BARQUAT® 4250, BARQUAT® 4280, BARQUAT® 4250Z, BARQUAT® 4280Z, BTC® 471, BTC® 2125, or BTC® 2125M each described as being a composition based on alkyldimethylbenzyl ammonium chloride and/or alkyldimethylethylbenzyl ammonium chloride; BARQUAT®  
20 MS-100 or BTC® 324-P-100 each described as being based on myristyldimethylbenzyl ammonium chloride; HYAMINE® 2389 described as being based on methyldodecylbenzyl ammonium chloride and/or methyldodecylxylene-bis-trimethyl ammonium chloride; HYAMINE® 1622 described  
25 as being an aqueous solution of benzethonium chloride; as well as BARQUAT® 1552 or BTC® 776 described as being based on alkyl dimethyl benzyl ammonium chloride and/or dialkyl methyl benzyl ammonium chloride, BARQUAT® 50-MAB described as being based on  
30 alkyldimethylethyl ammonium bromide and LONZABAC®-12.100 described as being based on an alkyl tertiary amine. (Each of these recited materials are presently

commercially available from Lonza, Inc., Fairlawn, NJ and/or from Stepan Co., Northfield IL)

The germidical constituent may be present in any effective amount, but generally need not be present in amounts in excess of about 10%wt. based on the total weight of the composition. The preferred germicidal cationic surfactant(s) may be present in the concentrated liquid disinfectant compositions in amounts of from about 0.001 % by weight to up to about 10% by weight, very preferably about 0.01-8% by weight, more preferably in amount of between 0.5-6 % by weight, and most preferably from 2 - 4% by weight.. It is particularly advantageous that the preferred germicidal cationic surfactant(s) are present in amounts of at least 200 parts per million (ppm), preferably in amounts of 200 - 700 ppm, more preferably in amounts of from 250 - 500 ppm, and very especially in amount of from 300 - 500ppm.

The inventive compositions necessarily include (b) at least one alkoxylated quaternary ammonium compound. This alkoxylated quaternary ammonium compound are selected to differ from the quaternary ammonium compounds discussed above, when the latter are present. These b) alkoxylated quaternary ammonium compound include materials which are per-se, known to the art.

Particularly useful alkoxylated quaternary ammonium compounds include those which may be represented by the general structure:



5

wherein: R is a C<sub>8</sub>-C<sub>24</sub> alkyl group;  
 z + y are individually integer values  
 of from 1 - 14, inclusive, but are most  
 preferably selected to that z + y = 2 -  
 15, and in certain particularly  
 preferred embodiments z + y = 2 or 15;  
 X may be any counterion, but is  
 desirably Cl or NO<sub>3</sub>.

Exemplary materials include ETHOQUAD 18/12  
 described to be octadecylmethyl [ethoxylated (2)]-  
 ammonium chloride; ETHOQUAD 18/25 described to be  
 octadecyl methyl [ethoxylated (15)] ammonium chloride,  
 ETHOQUAD C/25 described to be coco methyl [ethoxylated  
 (15)] ammonium chloride, ETHOQUAD C/12 described to be  
 coco methyl [ethoxylated (2)] ammonium chloride,  
 ETHOQUAD C/12 Nitrate described to be coco methyl  
 [ethoxylated (2)] ammonium nitrate, ETHOQUAD O/25  
 described to be oleyl methyl [ethoxylated (15)]  
 ammonium chloride, ETHOQUAD O/12 described to be oleyl  
 methyl [ethoxylated (2)] ammonium chloride, as well as  
 ETHOQUAD T/12 described to be tallow alkyl methyl  
 [ethoxylated (2)] ammonium chloride.

Further exemplary materials include Q-18-15 described to be octadecyl poly(15)oxyethylene methyl ammonium chloride and Q-C-15 described to be coco poly(15)oxyethylene methyl ammonium chloride (both of which are available from Tomah Inc.), as well as VARIQUAT K-1215, a methyl bis-(polyethoxy ethanol) coco ammonium chloride, with an 15 ethoxy groups; ADOGEN 66, an ethyl bis-(polyethoxy ethanol) tallow ammonium chloride, having 15 ethoxy groups, VARISOFT 5TD, an ethoxylated di (C<sub>12</sub>-C<sub>18</sub>) alkyl methyl ammonium chloride, with 5 ethoxy groups, REWOQUAT CPEM, a coco pentaethoxy methyl ammonium methosulfate, with 5 ethoxy groups.

An exemplary and particularly preferred alkoxylated quaternary ammonium compound is commercially available as ETHOQUAD 18/25 (ex. AKZO Corp.). Other particularly preferred compounds are described with reference to the Examples described below.

According to the first aspect and second aspects of the invention, the compositions necessarily include (c) one or more surfactants which provide a further deterative benefit to the compositions, ( but it is to be understood that according to further certain specific particularly embodiments these one or more surfactants are specifically absent.)

Useful surfactants which provide a further deterative benefit which may be present in the inventive compositions include deterative surfactants particularly selected from carboxylate, nonionic, cationic and amphoteric surfactants.

Suitable nonionic surfactants include, inter alia, condensation products of alkylene oxide groups with an organic hydrophobic compound, such as an aliphatic compound or with an alkyl aromatic compound.

5 The nonionic synthetic organic detergents generally are the condensation products of an organic aliphatic or alkyl aromatic hydrophobic compound and hydrophilic ethylene oxide groups. Practically any hydrophobic compound having a carboxy, hydroxy, amido, or amino

10 group with a free hydrogen attached to the nitrogen can be condensed with ethylene oxide or with the polyhydration product thereof, polyethylene glycol, to form a water soluble nonionic detergent. Further, the length of the polyethenoxy hydrophobic and hydrophilic

15 elements may be varied to adjust these properties.

One example of such a nonionic surfactant is the condensation product of one mole of an alkyl phenol having an alkyl group containing from 6 to 12 carbon atoms with from about 5 to 25 moles of an alkylene

20 oxide. Another example of such a nonionic surfactant is the condensation product of one mole of an aliphatic alcohol which may be a primary, secondary or tertiary alcohol having from 6 to 18 carbon atoms with from 1 to about 10 moles of alkylene oxide. Preferred

25 alkylene oxides are ethylene oxides or propylene oxides which may be present singly, or may be both present.

Preferred nonionic surfactants include primary and secondary linear and branched alcohol ethoxylates,

30 such as those based on C6 to C18 alcohols which further include an average of from 2 to 80 moles of ethoxylation per mol of alcohol. Particularly


preferred nonionic surfactants are C<sub>11</sub> linear primary alcohol ethoxylates averaging about 9 moles of ethylene oxide per mole of alcohol. These surfactants are available, for example, under the commercial name of Neodol 1-9, (from Shell Chemical Company, Houston, TX) , or in the Genapol® series of linear alcohol ethoxylates, particularly Genapol® 26-L-60 or Genapol® 26-L-80 (from Clariant Corp., Charlotte, NC). A further class of nonionic surfactants which are advantageously present in the inventive compositions are those presently marketed under the Genapol® tradename.

A further particularly useful and preferred alcohol ethoxylate is Genapol® UD-079 which is described to be a C<sub>11</sub> linear alcohol condensed with 7 moles of ethylene oxide to form a nonionic surfactant.


It is to be understood that other nonionic surfactants other than those described above may also be used. By way of illustration, and not by way of limitation, examples include secondary C<sub>12</sub> to C<sub>15</sub> alcohol ethoxylates, including those which have from about 3 to about 10 moles of ethoxylation. Such are available in the Tergitol® series of nonionic surfactants (Union Carbide Corp., Danbury, CT), particularly those in the Tergitol® "15-S-" series. Further exemplary nonionic surfactants include linear primary C<sub>11</sub> to C<sub>15</sub> alcohol ethoxylates, including those which have from about 3 to about 10 moles of ethoxylation. Such are available in the Neodol® series of nonionic surfactants (Shell Chemical Co.)

A further class of nonionic surfactants which may find use in the present inventive compositions include

ethoxylated octyl and nonyl phenols include those having one of the following general structural formulas:



5 or,



in which the  $\text{C}_9\text{H}_{19}$  group in the latter formula is a mixture of branched chained isomers, and  $x$  indicates an average number of ethoxy units in the side chain.

10 Particularly suitable non-ionic ethoxylated octyl and nonyl phenols include those having from about 7 to about 13 ethoxy units. Such compounds are commercially available under the trade name Triton® X (Union Carbide, Danbury CT), as well as under the  
15 tradename Igepal® (Rhône-Poulenc, Princeton, NJ). One exemplary and particularly preferred nonylphenol ethoxylate is Igepal® CO-630.

One useful class of surfactants include amine oxide compounds. Exemplary useful amine oxide  
20 compounds may be defined as one or more of the following of the four general classes:

(1) Alkyl di (lower alkyl) amine oxides in which the alkyl group has about 6-24, and preferably 8-18 carbon atoms, and can be straight or branched chain,  
25 saturated or unsaturated. The lower alkyl groups include between 1 and 7 carbon atoms, but preferably each include 1 - 3 carbon atoms.. Examples include



octyl dimethyl amine oxide, lauryl dimethyl amine oxide, myristyl dimethyl amine oxide, and those in which the alkyl group is a mixture of different amine oxides, such as dimethyl cocoamine oxide, dimethyl  
5 (hydrogenated tallow) amine oxide, and myristyl/palmityl dimethyl amine oxide;


(2) Alkyl di (hydroxy lower alkyl) amine oxides in which the alkyl group has about 6-22, and preferably 8-18 carbon atoms, and can be straight or  
10 branched chain, saturated or unsaturated. Examples include bis-(2-hydroxyethyl) cocoamine oxide, bis-(2-hydroxyethyl) tallowamine oxide; and bis-(2-hydroxyethyl) stearylamine oxide;

(3) Alkylamidopropyl di(lower alkyl) amine oxides in which the alkyl group has about 10-20, and preferably 12-16 carbon atoms, and can be straight or  
15 branched chain, saturated or unsaturated. Examples are cocoamidopropyl dimethyl amine oxide and tallowamidopropyl dimethyl amine oxide; and

20 (4) Alkylmorpholine oxides in which the alkyl group has about 10-20, and preferably 12-16 carbon atoms, and can be straight or branched chain, saturated or unsaturated.

While these amine oxides recited above may be  
25 used, preferred are amine oxides which may be represented by the following structural representation:

16



wherein

each  $\text{R}_1$  independently is a straight chained  $\text{C}_1\text{-C}_4$  alkyl group, preferably both  $\text{R}_1$  are methyl groups; and,

$\text{R}_2$  is a straight chained  $\text{C}_6\text{-C}_{22}$  alkyl group, preferably is  $\text{C}_6\text{-C}_{16}$  alkyl group, most preferably is a  $\text{C}_{8-10}$  alkyl group, especially a  $\text{C}_8$  alkyl group;

Each of the alkyl groups may be linear or branched, but most preferably are linear. Most preferably the amine oxide constituent is lauryl dimethyl amine oxide. Technical grade mixtures of two or more amine oxides may be used, wherein amine oxides of varying chains of the  $\text{R}_2$  group are present. Preferably, the amine oxides used in the present invention include  $\text{R}_2$  groups which comprise at least 50%wt., preferably at least 75%wt. of  $\text{C}_8$  alkyl group.

Exemplary and preferred amine oxide compounds include N-alkyl dimethyl amine oxides, particularly octyl dimethyl amine oxides as well as lauryl dimethyl amine oxide. These amine oxide compounds are available as surfactants from McIntyre Group Ltd. under the name Mackamine® C-8 which is described as a 40% by weight active solution of octyl dimethyl amine oxide, as well as from Stepan Co., under the tradename Ammonyx® LO which is described to be as a 30%wt. active solution of lauryl dimethyl amine oxide.

A further class of materials surfactants which may be advantageously included in the inventive compositions are alkoxy block copolymers, and in particular, compounds based on ethoxy/propoxy block copolymers. Polymeric alkylene oxide block copolymers include nonionic surfactants in which the major portion of the molecule is made up of block polymeric C<sub>2</sub>-C<sub>4</sub> alkylene oxides. Such nonionic surfactants, while preferably built up from an alkylene oxide chain starting group, and can have as a starting nucleus almost any active hydrogen containing group including, without limitation, amides, phenols, thiols and secondary alcohols.

One group of such useful nonionic surfactants containing the characteristic alkylene oxide blocks are those which may be generally represented by the formula (A):



where EO represents ethylene oxide,  
PO represents propylene oxide,  
y equals at least 15,  
(EO)<sub>x+z</sub> equals 20 to 50% of the total weight of said compounds, and,  
the total molecular weight is preferably in the range of about 2000 to 15,000.

Another group of nonionic surfactants appropriate for use in the new compositions can be represented by the formula (B):



wherein R is an alkyl, aryl or aralkyl group, where the R group contains 1 to 20 carbon atoms, the weight

percent of EO is within the range of 0 to 45% in one of the blocks a, b, and within the range of 60 to 100% in the other of the blocks a, b, and the total number of moles of combined EO and PO is in the range of 6 to 125 moles, with 1 to 50 moles in the PO rich block and 5 to 100 moles in the EO rich block.

Further nonionic surfactants which in general are encompassed by Formula B include butoxy derivatives of propylene oxide/ethylene oxide block polymers having molecular weights within the range of about 2000-5000.

Still further useful nonionic surfactants containing polymeric butoxy (BO) groups can be represented by formula (C) as follows:


$$\text{RO}-(\text{BO})_n(\text{EO})_x-\text{H} \quad (\text{C})$$
wherein R is an alkyl group containing 1 to 20 carbon atoms,  
n is about 5-15 and x is about 5-15.

Also useful as the nonionic block copolymer surfactants, which also include polymeric butoxy groups, are those which may be represented by the following formula(D):

$$\text{HO}-(\text{EO})_x(\text{BO})_n(\text{EO})_y-\text{H} \quad (\text{D})$$
wherein n is about 5-15, preferably about 15,  
x is about 5-15, preferably about 15,  
and  
y is about 5-15, preferably about 15.

Still further useful nonionic block copolymer surfactants include ethoxylated derivatives of

propoxylated ethylene diamine, which may be represented by the following formula:



where (EO) represents ethoxy,

5 (PO) represents propoxy,

the amount of (PO)<sub>x</sub> is such as to provide a molecular weight prior to ethoxylation of about 300 to 7500, and the amount of (EO)<sub>y</sub> is such as to provide about 20% to 90% of the total weight of said compound.


10 Of these, the most preferred are those which are represented by formula (A) above; specific examples of which include those materials presently commercially available under the tradename "Pluronic®", and in particular the Pluronic® F series, Pluronic® L series, 15 Pluronic® P series, as well as in the Pluronic® R series, each of which are generally described to be block copolymers of propylene oxide and ethylene oxide. Generally those of the Pluronic® L series and the Pluronic® R series are preferred as these are 20 supplied in liquid form by the manufacturer and are readily formulated into the present inventive compositions. These are also available in a wide range of HLB values, and those having HLB values in the range of 1.0 - 23.0 may be used, although those 25 with intermediate HLB values such as from about 12.0 - 18.0 are found to be particularly advantageous. These materials are presently commercially available from BASF AG (Ludwigshafen, Germany) as well as from BASF Corp. (Mt. Olive Township, New Jersey).

A further class of surfactants which may be advantageously included in the inventive compositions are carboxylates, particularly one or more alkylpolyoxycarboxylates including  
 5 alkyletherpolyoxycarboxylates, or alkylarylpolycarboxylates. Exemplary alkylpolyoxycarboxylates and alkylarylpolycarboxylates include alkyl- and alkylaryl-carboxylates which include those which may be represented by the general  
 10 formula:



wherein R is a straight or branched hydrocarbon chain containing from about 9 to 21 carbon atoms, and which may also include an aromatic ring, especially a  
 15 phenyl group as part of the hydrocarbon chain, and M is a metal or ammonium ion.

Further examples of particularly useful carboxylate surfactants include compounds according to the formula:



20

where:

R is a C<sub>4</sub>-C<sub>22</sub> linear or branched alkyl group which may optionally include at least one aryl group, preferably C<sub>8</sub>-C<sub>15</sub> linear or  
 25 branched alkyl group which may include at least one aryl group, and yet more preferably a C<sub>12-15</sub> linear or branched alkyl group which may include at least one aryl group;  
 x is an integer from 1 to 24,


y is 0 or 1,

R<sub>1</sub>, R<sub>2</sub> and R<sub>3</sub> is a group selected from H, lower alkyl radicals including methyl and ethyl radicals, carboxylate radicals including acetate and propionate radicals, succinate radicals, hydroxysuccinate radicals, or mixtures thereof wherein at least one R<sub>1</sub>, R<sub>2</sub> or R<sub>3</sub> is a carboxylate radical; and,

M<sup>+</sup> is a counterion including an alkali metal counterion (i.e., sodium, potassium) or ammonium counterion.

Free acid forms of the alkylethercarboxylate compounds noted above may also be used.

Examples of such presently available commercial preparations include SURFINE WLG (Finetex Inc., Elmwood Park NJ), SANDOPAN DTC (Clariant Chem.Co., Charlotte NC) in salt forms, and in free acid forms include those marketed under the tradename NEODOX (Shell Chemical Co., Houston TX). One particularly preferred carboxylate is one which is represented by the formula:



Such a material is presently commercially available under the tradename Emcol®, and specifically as Emcol® CNP-110.

Other useful exemplary nonionic block copolymers based on a polymeric ethoxy/propoxy units which may also be used include those presently commercially

available in the Poly-Tergent® E, and Poly-Tergent® P series of materials from Olin Chemicals Corp., (Stamford CT). These are described to be nonionic surfactants based on ethoxy/propoxy block copolymers,  
5 conveniently available in a liquid form from its supplier.

It is to be understood that these nonionic surfactants based on polymeric alkylene oxide block copolymers may be used singly or in mixtures of two or  
10 more such compounds.

Amphoteric surfactants, also known as zwitterionic surfactants, contain both cationic and anionic hydrophilic groups on the same molecule at a relatively wide range of pHs. The typical cationic  
15 group is a quaternary ammonium group, although other positively charged groups, like sulfonium groups, can also be used. The typical anionic hydrophilic groups are carboxylates and sulfonates, although other groups like sulfates, etc., can be used. Amphoteric  
20 surfactants also include betaine and sulphobetaine surfactants, derivatives thereof, and mixtures thereof wherein the molecule contains both basic and acidic groups which form an inner salt giving the molecule both cationic and anionic hydrophilic groups over a  
25 broad range of pH values, as well as mono- and diacetates, glycines, imidazolines and their derivatives, isethionates, mono- and dipropionates, hydroxy sultaines, and taurates.

When the compositions of the present invention  
30 contain one or more further deterative surfactants, these may be present in any amount which is found to provide a beneficial deterative effect. Generally,



these one or more further deterative surfactants do not comprise more than 12%wt. (on an actives weight basis) of the inventive compositions. When included such one or more further deterative surfactants are

5     advantageously present in an amount from 0.001 - 10%wt., preferably are present from 0.01 - 8%wt., but still more preferably are included in amounts of from 0.1 - 8%wt.

According to the first aspect and third aspects of the invention, the compositions necessarily include

10     (d) one or more organic solvents, ( but it is to be understood that according to further certain specific particularly embodiments these one or more surfactants are specifically absent.)

15     Exemplary organic solvents which may be included in the inventive compositions include those which are at least partially water-miscible such as alcohols, water-miscible ethers (e.g. diethylene glycol diethylether, diethylene glycol dimethylether,

20     propylent glycol dimethylether), water-miscible glycol ether (e.g. propylene glycol monomethylether, propylene glycol mono ethylether, propylene glycol monopropylether, propylene glycol monobutylether,

25     ethylene glycol monobutylether, dipropylene glycol monomethylether, diethyleneglycol monobutylether), lower esters of monoalkylethers of ethyleneglycol or propylene glycol (e.g. propylene glycol monomethyl ether acetate) all commercially available such as from Union Carbide (Danbury, CT), Dow Chemical Co. (Midland,

30     MI) or Hoescht (Germany).: Mixtures of several organic solvents can also be used.

Preferred as solvents in this invention are the glycol ethers having the general structure  $R_a-O-R_b-OH$ , wherein  $R_a$  is an alkoxy of 1 to 20 carbon atoms, or aryloxy of at least 6 carbon atoms, and  $R_b$  is an ether condensate of propylene glycol and/or ethylene glycol having from one to ten glycol monomer units. Preferred are glycol ethers having one to five glycol monomer units. These are  $C_3-C_{20}$  glycol ethers. Examples of more preferred solvents include propylene glycol methyl ether, dipropylene glycol methyl ether, tripropylene glycol methyl ether, propylene glycol isobutyl ether, ethylene glycol methyl ether, ethylene glycol ethyl ether, ethylene glycol butyl ether, diethylene glycol phenyl ether, propylene glycol phenol ether, and mixtures thereof.

The compositions are largely aqueous in nature, and comprise as a further necessary constituent (e) water. Water is added to order to provide to 100% by weight of the compositions of the invention. The water may be tap water, but is preferably distilled and is most preferably deionized water. If the water is tap water, it is preferably substantially free of any undesirable impurities such as organics or inorganics, especially minerals salts which are present in hard water which may thus undesirably interfere with the operation of the constituents present in the aqueous compositions according to the invention.

As discussed previously, the inventive compositions may comprise one or more conventional optional additives. By way of non-limiting example, these include: pH adjusting agents and pH buffers

including organic and inorganic salts; non-aqueous solvents, perfumes, perfume carriers, optical brighteners, coloring agents such as dyes and pigments, opacifying agents, hydrotropes, antifoaming agents, viscosity modifying agents such as thickeners, enzymes, anti-spotting agents, anti-oxidants, anti-corrosion agents as well as others not specifically elucidated here. These ingredients may be present in any combinations and in any suitable amount that is sufficient for imparting the desired properties to the compositions. These one or more conventional additives, when present, should be present in minor amounts, preferably in total comprise less than about 5% by weight (on an active weight basis) of the compositions, and desirably less than about 3%wt.

Such materials described above are known to the art, including those described in *McCutcheon's Emulsifiers and Detergents (Vol.1)*, *McCutcheon's Functional Materials (Vol. 2)*, North American Edition, 1991; *Kirk-Othmer, Encyclopedia of Chemical Technology*, 3rd Ed., Vol. 22, the contents of which are herein incorporated by reference. For any particular composition, any optional constituents should be compatible with the other ingredients present.

The aqueous compositions according to the invention are desirably provided as a ready to use product which may be directly applied to a hard surface. Hard surfaces which are to be particularly denoted are lavatory fixtures, lavatory appliances (toilets, bidets, shower stalls, bathtubs and bathing appliances), wall and flooring surfaces especially

those which include refractory materials and the like. Further hard surfaces which are particularly denoted are those associated with kitchen environments and other environments associated with food preparation.

5 Hard surfaces which are those associated with hospital environments, medical laboratories and medical treatment environments. Such hard surfaces described above are to be understood as being recited by way of illustration and not be way of limitation.

10 The composition provided according to the invention can be desirably provided as a ready to use product in a manually operated spray dispensing container, or may be supplied in aerosolized product wherein it is discharged from a pressurized aerosol  
15 container. Known art propellants such as liquid propellants based on chlorofluorocarbons or propellants of the non-liquid form, i.e., pressurized gases, including carbon dioxide, air, nitrogen, as well as others, may be used, even though it is realized that  
20 the former chlorofluorocarbons are not generally further used due to environmental considerations.

The composition according to the invention is ideally suited for use in a consumer "spray and wipe" application. In such an application, the consumer  
25 generally applies an effective amount of the cleaning composition using the pump and within a few moments thereafter, wipes off the treated area with a rag, towel, or sponge, usually a disposable paper towel or sponge. In certain applications, however, especially  
30 where undesirable stain deposits are heavy, the cleaning composition according to the invention may be left on the stained area until it has effectively

loosened the stain deposits after which it may then be wiped off, rinsed off, or otherwise removed. For particularly heavy deposits of such undesired stains, multiple applications may also be used. Where  
5 thorough disinfection is a primary consideration, it may be desired to apply the inventive compositions to the hard surface being treated and to permit the composition to remain on the hard surface for several minutes (2-10 min.) prior to rinsing or wiping the  
10 composition from the hard surface. It is also contemplated that the inventive compositions be applied to a hard surface without subsequently wiping or rinsing the treated hard surface.

In a yet a further embodiment, the product  
15 according to the invention may be formulated so that it may be useful in conjunction with a "aerosol" type product wherein it is discharged from a pressurized aerosol container. Known art propellants such as liquid propellants based on chlorofluorocarbons or  
20 propellants of the non-liquid form, i.e., pressurized gases, including carbon dioxide, air, nitrogen, as well as others, may be used, even though it is realized that the former chlorofluorocarbons are not generally further used due to environmental  
25 considerations. In such an application, the cleaning composition is dispensed by activating the release nozzle of said aerosol type container onto the stain and/or stain area, and in accordance with a manner as above-described a stain is treated and removed.

30 Whereas the compositions of the present invention are intended to be used in the types of liquid forms described, nothing in this specification shall be

understood as to limit the use of the composition according to the invention with a further amount of water to form a cleaning solution therefrom. In such a proposed diluted cleaning solution, the greater the proportion of water added to form said cleaning  
5 dilution will, the greater may be the reduction of the rate and/or efficacy of the thus formed cleaning solution. Accordingly, longer residence times upon the stain to effect their loosening and/or the useage  
10 of greater amounts may be necessitated. Conversely, nothing in the specification shall be also understood to limit the forming of a "super-concentrated" cleaning composition based upon the composition described above. Such a super-concentrated ingredient  
15 composition is essentially the same as the cleaning compositions described above except in that they include a lesser amount of water.

The following examples below illustrate exemplary and preferred formulations of the concentrate  
20 composition according to the instant invention. It is to be understood that these examples are presented by means of illustration only and that further useful formulations fall within the scope of this invention and the claims may be readily produced by one skilled  
25 in the art and not deviate from the scope and spirit of the invention.

Throughout this specification and in the accompanying claims, weight percents of any constituent are to be understood as the weight percent  
30 of the active portion of the referenced constituent, unless otherwise indicated.

Examples:

The following examples illustrate the formulation and performance of various compositions of the invention, as well as certain particularly preferred  
5       embodiments of the invention.

Exemplary formulations illustrating certain preferred embodiments of the inventive compositions and described in more detail in Table 1 below were formulated generally in accordance with the following  
10       protocol. The weight percentages indicated the "as supplied" weights of the named constituent.

Into a suitably sized vessel, a measured amount of water was provided after which the constituents were added in no specific or uniform sequence, which  
15       indicated that the order of addition of the constituents was not critical. All of the constituents were supplied at room temperature, and any remaining amount of water was added thereafter. Certain of the nonionic surfactants if gels at room  
20       temperature were first preheated to render them pourable liquids prior to addition and mixing. Mixing of the constituents was achieved by the use of a mechanical stirrer with a small diameter propeller at the end of its rotating shaft. Mixing, which  
25       generally lasted from 5 minutes to 120 minutes was maintained until the particular exemplary formulation appeared to be homogeneous. The exemplary compositions were readily pourable, and retained well mixed characteristics (i.e., stable mixtures) upon  
30       standing for extend periods. The compositions of the example formulations are listed on Table 1. Certain comparative examples are also described on Table 1;

these are identified as "C" followed by an integer number.

TABLE 1				
	C1	C2	Ex.1	Ex.2
BTC-8358 (80%)	3.6	--	3.6	3.6
ETHOQUAD 18/25 (95%)	--	2.12	0.68	2.12
di water	q.s.	q.s.	q.s.	q.s.

5 As is indicated, to all of the formulations of Table 1 was added sufficient deionized water in "quantum sufficient" to provide 100 parts by weight of a particular formulation.

10 The identity of the constituents of used to produce various formulations described herein are disclosed on Table 2, below, including the "actives" percentage of each where a constituent was not 100%wt. "actives".

TABLE 2	
BTC-8358 (80%)	alkyl dimethyl benzyl ammonium chloride (80%wt. actives) from Stepan Co.
ETHOQUAD 18/25 (95%)	octodecyl methyl [ethoxylated (15)] ammonium chloride (95%wt. actives) from Akzo-Nobel Co.
di water	deionized water

15

### Cleaning Efficacy

20 Certain of the compositions indicated above were diluted with water at a respective weight ratio of composition:water of 1:64. These diluted compositions were then subjected to the protocol of ASTM D-4488-89 Annex A5 for particulate soil, which evaluated the efficacy of the cleaning compositions on vinyl tile



samples. The soil applied was a particulate soil sample containing natural humus, paraffin oil, used crankcase motor oil, Portland cement, silica, lampblack carbon, iron oxide, bandy black clay, stearic acid, and oleic acid. produced according to the protocol. Each of the soiled test vinyl tile samples were placed into the apparatus and the center of each tile was wetted with a 20 milliliter sample of a test formulation and allowed to stand for 1 minute. When approximately 30 seconds had elapsed, a further 50 milliliter sample was applied to the sponge (water dampened, then wrung to remove excess water) of a Gardner Abrasion Tester apparatus. Thereafter the apparatus was cycled 2, 4, 6, 8 and 10 times, which provided, respectively 4, 8, 12, 16 and 20 strokes of the sponge across the face of each of the vinyl test tiles. The reflectance values of the cleaned samples at 2, 4, 6, 8 and 10 cycles were evaluated utilizing a Minolta Chroma Meter CF-110, with Data Processor DP-100, which evaluated spectrophotomic characteristics of the sample. These readings are reported on Table 3, following.

Table 3					
Formu- ation	2 cycles	4 cycles	6 cycles	8 cycles	10 cycles
C1	8.8	10.9	32.6	35.6	45.0
C2	16.3	23.0	23.5	29.9	23.6
C3	25.9	31.4	56.5	60.6	66.8
Ex.1	57.4	62.8	61.7	66.8	65.4
Ex.2	70.9	81.6	72.3	73.3	73.1

Comparative "C3" was a commercially available cleaning and disinfecting composition, LYSOL Disinfectant Cleaner, "Country Scent" (ex Reckitt &

Colman Inc., Wayne, NJ) which was diluted and tested in the manner described above. The composition of C3 was used as a benchmark for cleaning performance.

With respect to the results reported on Table 3 a value of "100" is indicative of a white (unsoiled) background, and a "0" value is indicative of a black background. As can be seen from the results of Table 3, the cleaning efficacy of the composition according to the invention provided superior results or were on parity with those of commercially available cleaning products. Surprisingly as can be gleaned from the results of C1, C2 in comparison with the results for Ex.1 and Ex.2 the inventive compositions provide unexpectedly superior cleaning particularly at 2, 4 and 6 cycles of the test. Such is unexpected, and is suggestive of a synergistic effect.

#### Antimicrobial Efficacy

The following inventive composition (identified as Ex. 3 below)

	Ex. 3
BTC-8358 (80%)	0.05625
ETHOQUAD 18/25 (95%)	0.05263
di water	q.s.

was evaluated for antimicrobial activity using the Biomek® 2000 Laboratory Automation Workstation together with the BioWorks Operating System (available from Beckman Coulter Inc., Fullerton, CA). The organism tested was Staphylococcus aureus at a concentration of 9 logs. The Biomek simulates a microbial reduction suspension test. One part of organism suspension (Staphylococcus aureus) is added to 9 parts of Ex. 3 in an appropriate container.

Deionized water (DI H<sub>2</sub>O) was used a control. The organism and sample are then mixed thoroughly for 15 seconds. Serial tenfold dilutions are carried out in a neutralizing broth. The diluted samples are then  
5 incubated for 24-48 hours at 35-37°C. Thereafter, surviving organisms are quantified and log reduction, as a measurement of organism survivors are calculated as follows:

Log Reduction = (Log Survivors/DI H<sub>2</sub>O Control) - (Log  
10 Survivors/Sample)

Ex. 3 had a log reduction of 4.8.

As may be seen from the results indicated above, the compositions according to the invention provide excellent cleaning benefits to hard surfaces,  
15 including hard surfaces with difficult to remove stains notwithstanding the low solids content of the inventive compositions. These advantages are further supplemented by the excellent antimicrobial efficacy of these compositions against known bacteria commonly  
20 found in bathroom, kitchen and other such advantages clearly illustrate the superior characteristics of the compositions, the cleaning and antimicrobial benefits attending its use which is not before known to the art.

25 While the invention is susceptible of various modifications and alternative forms, it is to be understood that specific embodiments thereof have been shown by way of example in the drawings which are not intended to limit the invention to the particular  
30 forms disclosed; on the contrary the intention is to cover all modifications, equivalents and alternatives

falling within the scope and spirit of the invention  
as expressed in the appended claims.


5

## Claims:

1. A hard surface cleaning and disinfecting composition which comprises:
  - 5 (a) a germicidal constituent which provides a primary sanitizing benefit, preferably a quaternary ammonium compound;
  - (b) an alkoxylated quaternary ammonium compound;
  - 10 (c) optionally, one or more deterative surfactants particularly selected from carboxylate, nonionic, cationic and amphoteric surfactants;
  - (d) optionally, one or more organic solvents;
  - 15 (e) water.
2. A hard surface cleaning and disinfecting composition according to claim 1 which is characterized as being essentially free of (d) one  
20 or more organic solvents.
3. A hard surface cleaning and disinfecting composition according to claim 1 which is characterized as being essentially free of (c) one  
25 or more deterative surfactants.
4. A hard surface cleaning and disinfecting composition according to claim 1 which is characterized as being essentially free of both  
30 (c) one or more deterative surfactants and (d) one or more organic solvents.

5. A hard surface cleaning and disinfecting composition according to any preceeding claim wherein the (a) germicidal constituent which provides a primary sanitizing benefit is a quaternary ammonium compound.

6. A hard surface cleaning and disinfecting composition according to any preceeding claim wherein the (b) alkoxyated quaternary ammonium compound may be represented by the general structure:



wherein: R is a C<sub>8</sub>-C<sub>24</sub> alkyl group;

z + y are individually integer values of from 1 - 14, inclusive, but are most preferably selected to that z + y = 2 - 15, and in certain particularly preferred embodiments z + y = 2 or 15;

X may be any counterion, but is desirably Cl or NO<sub>3</sub>.

7. A hard surface cleaning and disinfecting composition according to claim 6 wherein R is a C<sub>8</sub>-C<sub>24</sub> alkyl group includes which includes an ethoxy group adjacent to the quaternary nitrogen atom.

8. A hard surface cleaning and disinfecting composition according to claim 1,3 or 5 wherein the (d) one or more organic solvents are selected from: alcohols, ethers glycol ethers, lower esters of monoalkylethers of ethyleneglycol or propylene glycol, as well as mixtures thereof.
9. A hard surface cleaning and disinfecting composition according to claim 1, 2, or 5 wherein the ( c) one or more deterative surfactants are selected from carboxylate, nonionic, cationic and amphoteric surfactants.
10. A hard surface cleaning and disinfecting composition substantially described with reference to the Examples.

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/GB 00/00464

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C11D3/48 C11D1/62 C11D1/645 C11D3/20 A01N33/12

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C11D A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DE 196 15 286 A (HENKEL KGAA) 23 October 1997 (1997-10-23) the whole document	1,3,8-10
A		2,4
X	GB 2 320 030 A (RECKITT & COLMAN INC) 10 June 1998 (1998-06-10) page 11, line 23 -page 12, line 10 examples 1-11,23,24,26,27,29,31-38,56-65	1,2,5,9, 10
X	US 4 895 667 A (FOX PRISCILLA S ET AL) 23 January 1990 (1990-01-23) column 2, line 21-64 column 3, line 1-31 examples I,III-VIII	1-6
	-/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "A" document member of the same patent family

Date of the actual completion of the international search

28 April 2000

Date of mailing of the international search report

15/05/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Bertran Nadal, J



# INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 00/00464

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 111 965 A (PROCTER & GAMBLE) 27 June 1984 (1984-06-27) example 2A page 54, line 20 -page 55, line 6	1,5,8,9
A	US 5 629 280 A (TARASCHI FREDERIC A ET AL) 13 May 1997 (1997-05-13) the whole document	1-5,8-10
A	EP 0 478 445 A (PETERS) 1 April 1992 (1992-04-01) page 2, line 57 -page 4, line 57	1,5,8-10

# INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/GB 00/00464

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 19615286 A	23-10-1997	NONE	
GB 2320030 A	10-06-1998	AU 5082998 A CN 1239864 A EP 0946091 A WO 9824314 A ZA 9710895 A	29-06-1998 29-12-1999 06-10-1999 11-06-1998 17-07-1998
US 4895667 A	23-01-1990	NONE	
EP 0111965 A	27-06-1984	US 4551506 A US 4664848 A US 4659802 A AT 44978 T AU 576744 B AU 2280383 A CA 1213275 A DE 3380259 D DK 598383 A EG 17033 A FI 834747 A,B, GB 2133028 A,B GB 2168708 A,B GB 2171710 A,B GB 2171108 A,B GR 79156 A HK 57890 A HK 58190 A HK 74490 A HK 74690 A IE 56487 B JP 2090277 C JP 7100800 B JP 59166599 A KR 9201796 B MX 162904 B NO 834731 A,B, SG 35390 G SG 41090 G SG 41790 G SG 45090 G CA 1209140 A CA 1211446 A	05-11-1985 12-05-1987 21-04-1987 15-08-1989 08-09-1988 28-06-1984 28-10-1986 31-08-1989 24-06-1984 30-12-1991 24-06-1984 18-07-1984 25-06-1986 03-09-1986 20-08-1986 02-10-1984 10-08-1990 10-08-1990 28-09-1990 28-09-1990 14-08-1991 02-09-1996 01-11-1995 19-09-1984 02-03-1992 08-07-1991 25-06-1984 07-09-1990 03-08-1990 03-08-1990 17-08-1990 05-08-1986 16-09-1986
US 5629280 A	13-05-1997	GB 2304112 A AU 700516 B AU 6687096 A BR 9609984 A CA 2228427 A CN 1198772 A EP 0842248 A GB 2304728 A,B WO 9706230 A US 5939374 A	12-03-1997 07-01-1999 05-03-1997 12-01-1999 20-02-1997 11-11-1998 20-05-1998 26-03-1997 20-02-1997 17-08-1999
EP 0478445 A	01-04-1992	FR 2667220 A AT 161285 T CA 2052160 A	03-04-1992 15-01-1998 29-03-1992

# INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/GB 00/00464

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0478445 A		DE 69128447 D	29-01-1998
		DE 69128447 T	14-05-1998